

HHS Vaccine Clinic Anaphylaxis Management



- Location of kit in a yellow or blue bin within aftercare recovery areas, each bin contains more than 1 anaphylaxis kit, alongside oral Loratadine (Claritin) for hives post-vaccination
- Kit contents:
 - EPINEPHrine 1 mg/mL ampoule, filter needle for withdrawing drug from ampoule, IM needle for administration (23G 25mm SafetyGlide™), 1 mL syringe, chlorhexidine wipes
- Notify pharmacy if any medication is used, so that pharmacy can refill as necessary
- As per medical directive 49008:

Alpha and beta-adrenergic agonist	<p>Epinephrine HCL 1:1,000 (1 mg/mL) IM STAT.</p> <p>Administer 0.3 mg which is 0.3 mL intramuscular injection</p> <p>Dose may be repeated twice, administered 3 to 5 minutes apart</p>	Vaccine recipient observed or indicates Anaphylaxis or acute hypersensitivity reaction to administration of vaccine		<ul style="list-style-type: none"> • Epinephrine dosing may be repeated twice, administered 3 to 5 minutes apart • Call for assistance as per the clinic's escalation protocol including calling 911 • Transfer patient to an Emergency Department immediately
-----------------------------------	---	---	--	---

